

Example:

$$c = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$


$$\min c_1 x_1 + c_2 x_2$$

$$\text{s.t. } \boxed{-x_1 + x_2 \leq 1}$$

$$x_1 \geq 0$$

$$x_2 \geq 0$$

$$x_2 \leq -1$$



4 cases can occur:

i) there exists a unique opt. sol.

ii) there exist infinitely many opt. sol.

and the set of opt. sol. is either bounded or unbounded

iii) opt. sol. value is $-\infty$, no feasible solution is optimal

iv) feasible set is empty

Visualizing - problems in standard form

$$\min c^T x$$



$$\text{s.t. } Ax = b$$

$$x \geq 0$$

Example:

$$x_1 + x_2 + x_3 = 1$$

$$x_1, x_2, x_3 \geq 0$$



Chapter 2: The geometry of linear progr.

2.1 Polyhedra and convex sets

Def: Let $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$. Then the set

$$\{x \in \mathbb{R}^n \mid A \cdot x \geq b\}$$

is a polyhedron.

In particular, the set of feasible solutions to an LP is a polyhedron.

A set of the form $\{x \in \mathbb{R}^n \mid A \cdot x = b, x \geq 0\}$ is a polyhedron in standard form.


Def: A set $S \subseteq \mathbb{R}^n$ is bounded if there is a constant $K \in \mathbb{R}$ such that

$$\|x\|_{\infty} \leq K \quad \forall x \in S.$$

Def: Let $a \in \mathbb{R}^n \setminus \{0\}$ and $b \in \mathbb{R}$

(i) The set $\{x \in \mathbb{R}^n \mid a^T \cdot x = b\}$ is a hyperplane


(ii) The set $\{x \in \mathbb{R}^n \mid a^T \cdot x \geq b\}$ is a halfspace



$$\begin{aligned} a^T x &= a^T y = b \\ \Rightarrow a^T \cdot (x - y) &= 0 \end{aligned}$$

Def: $S \subseteq \mathbb{R}^n$ is convex, if

$$\lambda \cdot x + (1 - \lambda) \cdot y \in S \quad \forall x, y \in S, \lambda \in [0, 1]$$



Def: Let $x^1, \dots, x^k \in \mathbb{R}^n$ and $\lambda_1, \dots, \lambda_k \in \mathbb{R}$ with $\lambda_1, \dots, \lambda_k \geq 0$ and $\lambda_1 + \dots + \lambda_k = 1$



(i) The vector $\sum_{i=1}^k \lambda_i \cdot x^i$ is a convex combination of x^1, \dots, x^k .

(ii) The convex hull of x^1, \dots, x^k is



the set of all convex combinations of those vectors.



Theorem

(i) The intersection of convex sets is convex

(ii) Every polyhedron is convex.

(iii) A convex combination of a finite number of elements of a convex set also belongs to that set.

(iv) The convex hull of a finite number of vectors is a convex set.

Proof: (i) For $i \in I$ let S_i be a convex set

Let $x, y \in \bigcap_{i \in I} S_i$ and $\lambda \in [0, 1]$

S_i is convex $\Rightarrow \lambda \cdot x + (1-\lambda) \cdot y \in S_i \quad \forall i \in I$

$\Rightarrow \lambda \cdot x + (1-\lambda) \cdot y \in \bigcap_{i \in I} S_i$

(ii) $P = \{x \mid A \cdot x \geq b\}$

$x, y \in P, \lambda \in [0, 1]$

$$A \cdot (\lambda \cdot x + (1-\lambda) \cdot y) = \lambda \cdot \underbrace{Ax}_{\geq b} + (1-\lambda) \cdot \underbrace{Ay}_{\geq b} \geq b$$

(iii) Proof by induction over # elements. k .

$k=2$ by def.

$k \rightarrow k+1 \quad x^1, \dots, x^{k+1} \in S, \lambda_1, \dots, \lambda_{k+1} \geq 0 \quad \sum_{i=1}^{k+1} \lambda_i = 1$

$\sum_{i=1}^{k+1} \lambda_i \cdot x^i$ Assume w.l.o.g. $\lambda_{k+1} < 1$

$$= \lambda_{k+1} \cdot x^{k+1} + (1-\lambda_{k+1}) \cdot \underbrace{\sum_{i=1}^k \frac{\lambda_i}{(1-\lambda_{k+1})} \cdot x^i}_{\text{convex comb. of } x^1, \dots, x^k} \in S$$

$$(iv) \quad x = \sum_{i=1}^k \lambda_i \cdot x^i, \quad y = \sum_{i=1}^k \mu_i \cdot x^i \quad \lambda_i, \mu_i \geq 0 \quad \sum_i \lambda_i = \sum_i \mu_i = 1$$


$$\lambda \cdot x + (1-\lambda) \cdot y = \sum_{i=1}^k (\lambda \cdot \lambda_i + (1-\lambda) \cdot \mu_i) \cdot x^i$$

$$\sum_{i=1}^k (\lambda \cdot \lambda_i + (1-\lambda) \cdot \mu_i) = \lambda \cdot \underbrace{\sum_{i=1}^k \lambda_i}_{=1} + (1-\lambda) \cdot \underbrace{\sum_{i=1}^k \mu_i}_{=1} = 1 \quad \square$$

2.2. Extreme points, vertices, and basic feasible sol.

Def: Let P be a polyhedron in \mathbb{R}^n

(i) $x \in P$ is an extreme point of P if
 $x \neq \lambda \cdot y + (1-\lambda) \cdot z \quad \forall y, z \in P \setminus \{x\}, \lambda \in [0, 1]$



(ii) $x \in P$ is a vertex of P if there exists $c \in \mathbb{R}^n$ such that $c^T x < c^T y \quad \forall y \in P \setminus \{x\}$.

Def: Let $P \subseteq \mathbb{R}^n$ be a polyhedron defined by

$$a_i^T x \geq \delta_i \quad \text{for } i \in M_1$$

$$a_i^T \cdot x = \delta_i \quad \text{for } i \in M_2$$

$$a_i^T \cdot x \leq \delta_i \quad \text{for } i \in M_3$$

If a vector x^* satisfies $a_i^T \cdot x^* = \delta_i$ for some i then the corresp. constraint is active or binding at x^*

Theorem: Let $x^* \in \mathbb{R}^n$ and $I = \{i \mid a_i^T \cdot x^* = \delta_i\}$. The following are equivalent:

(i) There exist n vectors in $\{a_i \mid i \in I\}$ which are linearly independent.

(ii) The vectors in $\{a_i \mid i \in I\}$ span \mathbb{R}^n .

(iii) The system of equations $a_i^T \cdot x = \delta_i$ for $i \in I$ has a unique solution.