

Topology

Dr. Carsten Schultz

Axel Werner

Problem Set 3

Deadline: 12 Nov 2008

Exercise 16.

4 points

Let X and Y be quasi-compact spaces. Show that $X \times Y$ is again quasi-compact.
Hint: For a given open cover of $X \times Y$ and $x \in X$ show that there is a finite subcover of $\{x\} \times Y$. Then use Proposition 2.36 from the lecture and the quasi-compactness of X .

Exercise 17.

4 points

Show that for any index set J and path connected spaces X_j , $j \in J$, the product space $\prod_{j \in J} X_j$ is again path connected.

Remark: If we replace “path connected” by “connected”, then the statement is still true. For a finite index set J this has been proved in Proposition 2.14, for infinite J it is more complicated...

Exercise 18.

4 points

Let (X, d) be a metric space. Show that the topology on X induced by d is the initial topology with respect to the functions $f_y : X \rightarrow \mathbb{R}, x \mapsto d(x, y)$ for $y \in X$.

Exercise 19.

4 points

Let $C \subset [0, 1]$ be the Cantor set.

(a) Show that $C \times C$ is homeomorphic to C .

(b) What are the connected components of C ?

== == == == == == == ==
== == == == == == == ==

== == == == == == == ==
== == == == == == == ==

== == == == == == == ==
== == == == == == == ==

== == == == == == == ==
== == == == == == == ==

Exercise 20.**(Tutorial)**

With $I = [0, 1]$ show that I^I has a countable dense subset D .

Remark: This implies that the closure of D has more points than there are sequences in D !

Exercise 21.**(Tutorial)**

Let $S = (\{0, 1\}, \{\emptyset, \{1\}, \{0, 1\}\})$ be the Sierpiński space and (X, \mathcal{T}) some topological space.

(a) Show that \mathcal{T} is the initial topology with respect to the functions

$$\chi_O : X \rightarrow S, \chi_O(x) = \begin{cases} 1 & \text{if } x \in O \\ 0 & \text{otherwise} \end{cases}, \quad \text{for all } O \in \mathcal{T}$$



(b) What is the necessary condition for the function $\phi : X \rightarrow S^{\mathcal{T}}, \phi(x)(O) = \chi_O(x)$ to be an embedding?

Exercise 22.**(Tutorial)**

Let M be a set and for $a \in M$ define $\delta_a \in \{0, 1\}^M$ by

$$\delta_a(b) = \begin{cases} 1 & \text{if } a = b, \\ 0 & \text{otherwise.} \end{cases}$$

What is the closure of the set $\{\delta_a \mid a \in M\}$? Does it depend on M ?



<http://www.evilmadscientist.com/article.php/fractalcookies>