

## Differential Geometry II: Analysis and Geometry on Manifolds

### Exercise Sheet 8

(Parallel transport, Riemannian manifolds)

due 10.01.2017

#### Exercise 1

5 points

Let  $M \subset \mathbb{R}$  be an interval and consider the vector bundle  $E = M \times \mathbb{R}^k$ ,  $k \in \mathbb{N}$ , equipped with some connection  $\nabla$ . Show that  $(E, \nabla)$  is trivial.

#### Exercise 2

5 points

Let  $(M, g)$  be a Riemannian manifold and  $\tilde{g} = e^{2u}g$  for some smooth function  $u: M \rightarrow \mathbb{R}$ . Show that between the corresponding Levi-Civita connections the following relation holds:

$$\tilde{\nabla}_X Y = \nabla_X Y + du(X)Y + du(Y)X - g(X, Y)\text{grad } u.$$

Here  $\text{grad } u \in \Gamma(TM)$  is the vector field uniquely determined by the condition  $du(X) = g(\text{grad } u, X)$  for all  $X \in \Gamma(TM)$ .

#### Exercise 3

5 points

Let  $(M, \langle \cdot, \cdot \rangle)$  be a 2-dimensional Riemannian manifold,  $\nabla$  its Levi-Civita connection. Show that there is a function  $K \in \mathcal{C}^\infty(M)$  such that

$$R^\nabla(X, Y)Z = K(\langle Y, Z \rangle X - \langle X, Z \rangle Y), \text{ for all } X, Y, Z \in \Gamma(TM).$$