

Exercise Sheet 6 (26.11.19)

Due date: 03.12.19

- To get the Übungsschein (necessary condition for the oral exam) you need to collect 60% of the total sum of points in each half of the semester. Each exercise sheet has 20 points. The total number of sheets is 12-14.
- Please work in fixed groups of 2 or 3 students.
- Please justify each step of your computations. *Results without any explanation are not accepted.* Please write in a readable way. *Unreadable handwriting will not be corrected.* Feel free to write your answers either in English or in German.
- Please turn in your homework directly to me at the beginning of the Tutorial or leave it in my letter box (MA 701, Frau Jean Downes). *No homework will be accepted after the deadline has passed.*

Exercise 1

(4 pts)

Consider the following planar linear systems of ODEs:

$$\begin{cases} \dot{x}_1 = -2x_1, \\ \dot{x}_2 = -3x_2, \end{cases} \quad \begin{cases} \dot{x}_1 = -2x_1, \\ \dot{x}_2 = -2x_2. \end{cases}$$

1. Sketch the phase portraits of both systems.
2. Prove that the systems are topologically conjugate by constructing explicitly the map between the orbits.

Exercise 2

(2 pts)

Consider the following system of ODEs in \mathbb{R}^2 :

$$\begin{cases} \dot{x}_1 = x_1 (1 - x_1^2 - x_2^2) - x_2 (1 + x_1^2 + x_2^2), \\ \dot{x}_2 = x_1 (1 + x_1^2 + x_2^2) + x_2 (1 - x_1^2 - x_2^2). \end{cases}$$

1. Rewrite the system of ODEs using polar coordinates, $(x_1, x_2) := r(\cos \theta, \sin \theta)$, $r > 0$, $\theta \in [0, 2\pi)$.
2. Find a periodic solution.

TURN OVER!

Exercise 3(7 pts)

Consider the following system of ODEs in \mathbb{R}^2 :

$$\begin{cases} \dot{x}_1 = x_1^4 + x_1 x_2, \\ \dot{x}_2 = -2x_2 - x_1^2 + x_1 x_2^2. \end{cases} \quad (1)$$

1. Linearize (1) around the fixed point $(0,0)$. What can you say about the stability of $(0,0)$ on the basis of the Poincaré-Lyapunov Theorem?
2. Find the center (linear) space $E^c(0,0)$. Construct an approximation of the center manifold $W^c(0,0)$.
Hint: The center manifold is parametrized, in a neighborhood of $(0,0)$, by $x_2 = h(x_1)$ for some function h . An approximation of $W^c(0,0)$ is given by the series expansion – say up to $O(x_1^5)$ – of the function h around $x_1 = 0$.
3. Find the first three nonzero terms of the series expansion of the system obtained by reducing (1) on $W^c(0,0)$. What can you say now about the stability of the fixed point $(0,0)$ for system (1)?

Exercise 4(7 pts)

Consider the following system of ODEs in \mathbb{R}^3 :

$$\begin{cases} \dot{x}_1 = \frac{1}{2}(x_3 - x_1) - x_2 + 2x_2^3, \\ \dot{x}_2 = \frac{1}{2}(x_3 - x_1), \\ \dot{x}_3 = x_3 - x_2^2(x_1 + x_3). \end{cases} \quad (2)$$

1. Find all fixed points.
2. Linearize system (2) around the fixed point $(0,0,0)$ and discuss the stability of $(0,0,0)$. Write the general solution of the linearized system.
3. Prove that $M := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 - 2x_2 + x_3 = 0\}$ is an invariant manifold for (2).
4. Reduce system (2) on M by eliminating the variable x_1 , thus getting a first-order ODE for x_2 and x_3 . Then eliminate the variable x_3 , thus obtaining a second-order ODE for x_2 . Find an integral of motion of the resulting ODE.
Is this an integral of motion of the full system (2)?