Lamperti transform for multi-type CBI

Nicoletta Gabrielli
(based on a joint work with J. Teichmann)

University of Zürich
Department of Banking and Finance

nicoletta.gabrielli@bf.uzh.ch

October 23, 2014

Structure of the talk

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Branching processes

Let

$$(\Omega, (X_t)_{t\geq 0}, (\mathcal{F}_t^{\natural})_{t\geq 0}, (\mathbb{P}^x)_{x\in\mathbb{R}_{\geq 0}})$$

be a time homogeneous Markov process. The process X is said to be an branching process if it satisfies the following property:

Branching property

For any $t \geq 0$ and $x_1, x_2 \in \mathbb{R}_{\geq 0}$, the law of X_t under $\mathbb{P}^{x_1+x_2}$ is the same as the law of $X_t^{(1)} + X_t^{(2)}$, where each $X^{(i)}$ has the same distribution as X under $\mathbb{P}^{x_i}$, for $i = 1, 2$.

Fourier–Laplace transform characterization

- There exists a function $\Psi : \mathbb{R}_{\geq 0} \times \mathbb{C}_{\leq 0} \to \mathbb{C}$ such that

$$\mathbb{E}^x\left[e^{uX_t}\right] = e^{x\Psi(t,u)},$$

  for all $x \in \mathbb{R}_{\geq 0}$ and $(t, u) \in \mathbb{R}_{\geq 0} \times \mathbb{C}_{\leq 0}$.
- On the set $\mathcal{Q} = \mathbb{R}_{\geq 0} \times \mathbb{C}_{\leq 0}$, the function Ψ satisfies the equation

$$\partial_t \Psi(t, u) \quad = R(\Psi(t, u)), \quad \Psi(0, u) = u\,.$$

The branching mechanism

The function R has the following Lévy-Khintchine form

$$R(u) = \beta u + \frac{1}{2}u^2\alpha + \int_0^\infty \left(e^{u\xi} - 1 - u\xi \mathbb{1}_{\{|\xi|\leq 1\}}\right) M(d\xi)\,,$$

where $\beta \in \mathbb{R}$, $\alpha \geq 0$ and M is a Lévy measure with support in $\mathbb{R}_{\geq 0}$.

Lévy processes

A time homogeneous Markov process Z is a Lévy process if the following three conditions are satisfied:

L1) $Z_0 = 0$ $\mathbb{P}$-a.s.

L2) Z has independent and stationary increments, i.e. for all $n \in \mathbb{N}$ and $0 \leq t_0 < t_1 < \ldots < t_{n+1} < \infty$

(independence) the random variables $\{Z_{t_{j+1}} - Z_{t_j}\}_{j=0,\ldots,n}$ are independent,

(stationarity) the distribution of $Z_{t_{j+1}} - Z_{t_j}$ coincides with the distribution of $Z_{(t_{j+1}-t_j)}$,

L3) (stochastic continuity) for each $a > 0$ and $s \geq 0$,

$$\lim_{t \to s} \mathbb{P}(|Z_t - Z_s| > a) = 0\,.$$

Relation with infinitely divisible distributions

- If Z is a Lévy process, then, for any $t \geq 0$, the random variable Z_t is infinitely divisible.
- The Fourier transform of a Lévy process takes the form:

$$\mathbb{E}^0\left[e^{\langle u, Z_t\rangle}\right] = e^{t\eta(u)}, \quad u \in \mathrm{i}\mathbb{R}$$

$$\eta(u) = \beta u + \frac{1}{2}u^2\alpha + \int \left(e^{u\xi} - 1 - u\xi \mathbb{1}_{\{|\xi|\leq 1\}}\right) M(d\xi),$$

  where $\beta \in \mathbb{R}$, $\alpha \geq 0$ and M is a Lévy measure in $\mathbb{R}$.
- The Fourier transform can be extended in the complex domain and the resulting Fourier–Laplace transform is well defined in

$$\mathcal{U} := \{u \in \mathbb{C} \mid \eta(\mathcal{R}e(u)) < \infty\}.$$

Lamperti transform

Theorem [Lamperti, 1967]

Let Z be a Lévy process with no negative jumps with Lévy exponent R, i.e.

$$\mathbb{E}^0\left[e^{uZ_t}\right] = e^{tR(u)}, \qquad u \in \mathcal{U}\,.$$

Define, for $t \geq 0$

$$X_t = x + Z_{\theta_t \wedge \tau_0^-} \quad \theta_t := \inf\left\{s > 0 \mid \int_0^s \frac{dr}{Z_r} > t\right\}.$$

Then X is a CB process with branching mechanism R.

Lamperti transform

Theorem 2 in [Caballero et al., 2013]

Let Z be a Lévy process with no negative jumps with Lévy exponent R, i.e.

$$\mathbb{E}^0\left[e^{uZ_t}\right] = e^{tR(u)}, \qquad u \in \mathcal{U}\,.$$

The time–change equation

$$X_t = x + Z_{\int_0^t X_r dr}$$

admits a unique solution, which is a CB process with branching mechanism R.

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Add immigration

A CBI-process with branching mechanism R and immigration mechanism F is a Markov process Z taking values in $\mathbb{R}_{\geq 0}$ satisfying

- there exist functions $\phi : \mathbb{R}_{\geq 0} \times \mathcal{U} \to \mathbb{C}$ and $\Psi : \mathbb{R}_{\geq 0} \times \mathcal{U} \to \mathbb{C}$ such that
$$\mathbb{E}^x\left[e^{uX_t}\right] = e^{\phi(t,u)+x\Psi(t,u)},$$
for all $x \in \mathbb{R}_{\geq 0}$ and $(t,u) \in \mathbb{R}_{\geq 0} \times \mathcal{U}$.
- On the set $\mathcal{Q} = \mathbb{R}_{\geq 0} \times \mathcal{U}$, the functions ϕ and Ψ satisfy the following system :
$$\begin{aligned} \partial_t \phi(t,u) &= F(\Psi(t,u)), \quad \phi(0,u) = 0, \\ \partial_t \Psi(t,u) &= R(\Psi(t,u)), \quad \Psi(0,u) = u\,. \end{aligned}$$

Add immigration

The immigration mechanism

The function F has the following Lévy-Khintchine form

$$F(u) = bu + \int_0^\infty \left(e^{u\xi} - 1\right) m(d\xi)\,,$$

with $b \geq 0$ and m is a Lévy measure on $\mathbb{R}_{\geq 0}$ such that $\int (1 \wedge \xi) m(d\xi) < \infty$.

Lamperti transform for CBI

Theorem 2 in [Caballero et al., 2013]

Let $Z^{(1)}$ be a Lévy process with no negative jumps and $Z^{(0)}$ an independent subordinator such that

$$\mathbb{E}^0\left[e^{uZ_t^{(1)}}\right] = e^{tR(u)} \quad \text{and} \quad \mathbb{E}^0\left[e^{uZ_t^{(0)}}\right] = e^{tF(u)}, \qquad u \in \mathcal{U}.$$

The time–change equation

$$X_t = x + Z_t^{(0)} + Z^{(1)}_{\int_0^t X_r dr}$$

admits a unique solution, which is a CBI process with branching mechanism R and immigration mechanism F.

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Definition

Let

$$(\Omega, (X_t)_{t\geq 0}, (\mathcal{F}_t^{\natural})_{t\geq 0}, (\mathbb{P}^x)_{x\in\mathbb{R}^m_{\geq 0}})$$

be a time homogeneous Markov process. The process X is said to be a multi–type CBI if it satisfies the following property:

See [Duffie et al., 2003, Barczy et al., 2014]

There exist functions $\phi : \mathbb{R}_{\geq 0} \times \mathcal{U} \to \mathbb{C}$ and $\Psi : \mathbb{R}_{\geq 0} \times \mathcal{U} \to \mathbb{C}^m$ such that

$$\mathbb{E}^x\left[e^{\langle u, X_t\rangle}\right] = e^{\phi(t,u)+\langle x, \Psi(t,u)\rangle},$$

for all $x \in \mathbb{R}^m_{\geq 0}$ and $(t, u) \in \mathbb{R}_{\geq 0} \times \mathcal{U}$, with $\mathcal{U} = \mathbb{C}^m_{\leq 0}$.

Generalized Riccati equations

On the set $\mathcal{Q} = \mathbb{R}_{\geq 0} \times \mathcal{U}$, the functions ϕ and Ψ satisfy the following system of generalized Riccati equations:

$$\begin{aligned} \partial_t \phi(t,u) &= F(\Psi(t,u)), \quad \phi(0,u) = 0, \\ \partial_t \Psi(t,u) &= R(\Psi(t,u)), \quad \Psi(0,u) = u\,. \end{aligned}$$

Lévy–Khintchine form for the vector fields

The functions F and R_k, for each $k = 1, \ldots, m$, have the following Lévy-Khintchine form

$$F(u) = \langle b, u \rangle + \int_{\mathbb{R}^m_{\geq 0} \setminus \{0\}} \left(e^{\langle u, \xi \rangle} - 1 \right) m(d\xi),$$

$$R_k(u) = \langle \beta_k, u \rangle + \frac{1}{2} u_k^2 \alpha_k + \int_{\mathbb{R}^m_{\geq 0} \setminus \{0\}} \left(e^{\langle u, \xi \rangle} - 1 - \mathbf{u}_k \boldsymbol{\xi}_k \mathbb{1}_{\{|\xi| \leq 1\}} \right) M_k(d\xi).$$

Admissible parameters

The set of parameters satisfies the following restrictions

- $b, \beta_i \in \mathbb{R}^m$, $i = 1, \ldots, m$,
- $\alpha_i \geq 0$,
- m, M_i, $i = 1, \ldots, m$, Lévy measures.

drift	
$b \in \mathbb{R}^m_{\geq 0}$,	
$(\beta_i)_k \geq 0$,	for all $i = 1, \ldots, m$ and $k \neq i$,

jumps	
$\operatorname{supp} m \subseteq \mathbb{R}^m_{\geq 0}$,	and $\int (\lvert\xi\rvert \wedge 1)\, m(d\xi) < \infty$,
$\operatorname{supp} M_i \subseteq \mathbb{R}^m_{\geq 0}$,	for all $i = 1, \ldots, m$ and
	$\int \big((\lvert(\xi)_{-i}\rvert + \lvert\xi_i\rvert^2) \wedge 1\big) M_i(d\xi) < \infty$.

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Remarks

Remarks

- It is possible to define $m+1$ independent Lévy processes $Z^{(0)}, Z^{(1)}, \ldots, Z^{(m)}$ taking values in $\mathbb{R}^m$ with Lévy exponents $F, R_1, \ldots, R_m$.

Remarks

- It is possible to define $m+1$ independent Lévy processes $Z^{(0)}, Z^{(1)}, \ldots, Z^{(m)}$ taking values in $\mathbb{R}^m$ with Lévy exponents $F, R_1, \ldots, R_m$.
- In [Kallsen, 2006] it has been proved that the time change equation

$$X_t = x + Z_t^{(0)} + \sum_{k=1}^{m} Z^{(k)}_{\int_0^t X_s^{(k)} ds}, \quad t \geq 0\,, \qquad (*)$$

admits a weak solution, i.e. there exists a probability space containing two processes (X, Z) such that (*) holds in distribution.

Remarks

- It is possible to define $m+1$ independent Lévy processes $Z^{(0)}, Z^{(1)}, \ldots, Z^{(m)}$ taking values in $\mathbb{R}^m$ with Lévy exponents $F, R_1, \ldots, R_m$.
- In [Kallsen, 2006] it has been proved that the time change equation

$$X_t = x + Z_t^{(0)} + \sum_{k=1}^{m} Z^{(k)}_{\int_0^t X_s^{(k)} ds}, \quad t \geq 0\,, \qquad (*)$$

  admits a weak solution, i.e. there exists a probability space containing two processes (X, Z) such that (*) holds in distribution.
- Moreover X has the distribution of a multi–type CBI with immigration mechanism F and branching mechanism $(R_1, \ldots, R_m)$.

Remarks

- It is possible to define $m+1$ independent Lévy processes $Z^{(0)}, Z^{(1)}, \dots, Z^{(m)}$ taking values in $\mathbb{R}^m$ with Lévy exponents $F, R_1, \dots, R_m$.
- In [Kallsen, 2006] it has been proved that the time change equation

$$X_t = x + Z_t^{(0)} + \sum_{k=1}^{m} Z^{(k)}_{\int_0^t X_s^{(k)} ds}, \quad t \geq 0\,, \qquad (*)$$

  admits a weak solution, i.e. there exists a probability space containing two processes (X, Z) such that (*) holds in distribution.
- Moreover X has the distribution of a multi–type CBI with immigration mechanism F and branching mechanism $(R_1, \dots, R_m)$.

Does there exist a pathwise solution of (*)?

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Results for multi–type CB

Theorem [G. and Teichmann, 2014]

Let $Z^{(1)}, \ldots, Z^{(m)}$ be independent $\mathbb{R}^m$-valued Lévy processes with

$$\mathbb{E}^0\left[e^{\left\langle u, Z_t^{(k)}\right\rangle}\right] = e^{tR_k(u)}, \qquad u \in \mathcal{U}\,,$$

where each R_k is of LK form with triplets given by a set of admissible parameters. Then the time–change equation

$$X_t = x + \sum_{k=1}^{m} Z^{(k)}_{\int_0^t X_s^{(k)} ds} \qquad t \geq 0\,,$$

admits a unique solution, which is a multi–type CB process with respect to the **time–changed filtration**.

Multiparameter time–change filtration

- Define
  $Z = (Z_1^{(1)}, \dots, Z_m^{(1)}, \dots, Z_1^{(m)}, \dots, Z_m^{(m)}) =: (Z^{(1)}, \dots, Z^{(m^2)})$.
- For all $\underline{s} = (s_1, \dots, s_{m^2}) \in \mathbb{R}_{\geq 0}^{m^2}$
  $\mathcal{G}_{\underline{s}}^{\natural} := \sigma\left(\{Z_{t_h}^{(h)},\ t_h \leq s_h,\ \text{for } h = 1, \dots, m^2\}\right)$.
- Complete it by $\mathcal{G}_{\underline{s}} = \bigcap_{n \in \mathbb{N}} \mathcal{G}_{\underline{s}^{(n)} + \frac{1}{n}}^{\natural} \vee \sigma(\mathcal{N})$.

Definition

A random variable $\underline{\tau} = (\tau_1, \dots, \tau_{m^2}) \in \mathbb{R}_{\geq 0}^{m^2}$ is a $(\mathcal{G}_{\underline{s}})$-stopping time if

$$\{\underline{\tau} \leq \underline{s}\} := \{\tau_1 \leq s_1, \dots, \tau_{m^2} \leq s_{m^2}\} \in \mathcal{G}_{\underline{s}},\ \text{for all } \underline{s} \in \mathbb{R}_{\geq 0}^{m^2}.$$

If $\underline{\tau}$ is a stopping time,

$$\mathcal{G}_{\underline{\tau}} := \{B \in \mathcal{G} \mid B \cap \{\underline{\tau} \leq \underline{s}\} \in \mathcal{G}_{\underline{s}} \text{ for all } \underline{s} \in \mathbb{R}_{\geq 0}^{m^2}\}.$$

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

Extension to multi–type CBI

- Let F be an immigration mechanism and $R = (R_1, \dots, R_m)$ a branching mechanism.
- Let $Z^{(0)}$ Lévy process with exponent F and $Z^{(i)}$ Lévy process with exponent R_i.
- Define, for $k = 0, \dots, m$,

$$\overline{Z}^{(k)} := (\overline{Z}_0^{(k)}, \underbrace{\overline{Z}_1^{(k)}, \dots, \overline{Z}_m^{(k)}}_{m \text{ coordinates}}) := (0, \underbrace{Z^{(k)}}_{m \text{ coordinates}}).$$

- Given $y = (1, x)$ with $x \in \mathbb{R}^m_{\geq 0}$, the previous result gives pathwise existence of

$$Y_t = y + \sum_{k=0}^{m} \overline{Z}^{(k)}_{\int_0^t Y_s^{(i)} ds}.$$

- It holds $Y = (1, X)$ where X is a $CBI(F, R)$.

Outline

1. Introduction
   - Real valued CB and Lamperti transform
   - Extension to real valued CBI
2. What is a multi–type CBI
   - Definitions
   - Some additional results
3. Lamperti transform for multi–type CBI
   - Multi–type CB and their time–change representation
   - Extension to multi–type CBI
   - Sketch of the proof

The one dimensional case

Question: Given a Lévy process Z taking values in $\mathbb{R}$, is there a solution of

$$X_t = x + Z_{\int_0^t X_s ds} \qquad ?$$

- For the one dimensional case see also [Caballero et al., 2013].

An ODE point of view in $\mathbb{R}_{\geq 0}$

Introduce

$$\tau(t) := \int_0^t X_s ds \,.$$

Does there exist a solution $\tau \in \mathbb{R}_{\geq 0}$ of

$$\begin{cases} \dot{\tau}(t) = x + Z(\tau(t)) \\ \tau(0) = 0 \end{cases} \qquad ?$$

An ODE point of view in $\mathbb{R}^m_{\geq 0}$

Introduce

$$\begin{aligned} \mathcal{Z} : \mathbb{R}^m_{\geq 0} &\to \mathbb{R}^m \\ \underline{s} &\mapsto \sum_{i=1}^{m} Z^{(i)}(s_i)\,. \end{aligned}$$

Does there exist a solution $\underline{\tau} \in \mathbb{R}^m_{\geq 0}$ of

$$\begin{cases} \dot{\underline{\tau}}(t) = x + \mathcal{Z}(\underline{\tau}(t))\,, \\ \underline{\tau}(0) = 0\,. \end{cases} \qquad ?$$

Construction of the time–change process

Theorem [G. and Teichmann, 2014]

There exists a solution of

$$\begin{cases} \dot{\underline{\tau}}((t_0, \tau_0, x); t) = (x + \mathcal{Z})(\underline{\tau}((t_0, \tau_0, x); t)), \\ \underline{\tau}((t_0, \tau_0, x); t_0) = \tau_0 \,, \end{cases}$$

for $t \geq t_0$ and $\tau_0 \in \mathbb{R}^m_{\geq 0}$.

Construction of the time–change process

Theorem [G. and Teichmann, 2014]

There exists a solution of

$$\begin{cases} \dot{\underline{\tau}}((t_0, \tau_0, \mathbf{x}); t) = (\mathbf{x} + \mathcal{Z})(\underline{\tau}((t_0, \tau_0, \mathbf{x}); t)), \\ \underline{\tau}((t_0, \tau_0, \mathbf{x}); t_0) = \tau_0 \,, \end{cases}$$

for $t \geq t_0$ and $\tau_0 \in \mathbb{R}^m_{\geq 0}$.

Construction of the time–change process

Theorem [G. and Teichmann, 2014]

There exists a solution of

$$\begin{cases} \dot{\underline{\tau}}((\mathbf{t}_0, \boldsymbol{\tau}_0, x); t) = (x + \mathcal{Z})(\underline{\tau}((\mathbf{t}_0, \boldsymbol{\tau}_0, x); t)), \\ \underline{\tau}((\mathbf{t}_0, \boldsymbol{\tau}_0, x); \mathbf{t}_0) = \boldsymbol{\tau}_0 \,, \end{cases}$$

for $t \geq t_0$ and $\tau_0 \in \mathbb{R}^m_{\geq 0}$.

Decomposition of $\mathcal{Z}$

The Lévy–Itô decomposition together with the **canonical form** of the admissible parameters give

$$\begin{aligned} Z_t^{(i)} =& \beta_i t + \sigma_i B_t^{(i)} + \int_0^t \int \xi 1_{\{|\xi|>1\}} \mathcal{J}^{(i)}(d\xi, ds) \\ &+ \int_0^t \int \xi 1_{\{|\xi|\leq 1\}} (\mathcal{J}^{(i)}(d\xi, ds) - M_i(d\xi) ds) \end{aligned}$$

where $\sigma_i = \sqrt{\alpha_i}$, $B^{(i)}$ is a process in $\mathbb{R}^m$ which evolves only along the the i-th coordinate as Brownian motion and $\mathcal{J}^{(i)}$ is the jump measure of the process $Z^{(i)}$.

- Decompose

$$Z^{(i)} =: \widetilde{Z}^{(i)} + \overset{\infty}{Z}{}^{(i)}$$

where $\widetilde{Z}^{(i)}$ and $\overset{\infty}{Z}{}^{(i)}$ are two stochastic processes on $\mathbb{R}^m$ defined by

$$\widetilde{Z}_i^{(i)}(t) := (\beta_i)_i t + \sigma_i B^{(i)}(t) + \int_0^t \int \xi_i 1_{\{|\xi|>1\}} \mathcal{J}^{(i)}(d\xi, ds) + \int_0^t \int \xi_i 1_{\{|\xi|\leq 1\}} \widetilde{\mathcal{J}}^{(i)}(d\xi, ds)$$

$$\widetilde{Z}_k^{(i)}(t) := 0, \qquad \text{for } k \neq i\,,$$

$$\overset{\infty}{Z}{}^{(i)}(t) := \overset{\infty}{\beta}_i t + \int_0^t \int (\xi - \xi_i e_i) 1_{\{|\xi|>1\}} \mathcal{J}^{(i)}(d\xi, ds) + \int_0^t \int (\xi - \xi_i e_i) 1_{\{|\xi|\leq 1\}} \widetilde{\mathcal{J}}^{(i)}(d\xi, ds)\,.$$

where $\overset{\infty}{\beta}_i = \beta_i - e_i(\beta_i)_i$ and $\widetilde{\mathcal{J}}^{(i)}$ is the compensated jump measure.

Approximation of the jump part

- Introduce, for all $\underline{s} \in \mathbb{R}^m_{\geq 0}$,

$$\overset{\sim}{\mathcal{Z}}(\underline{s}) := \sum_{i=1}^{m} \tilde{Z}^{(i)}(s_i), \quad \overset{\infty}{\mathcal{Z}}(\underline{s}) := \sum_{i=1}^{m} \overset{\infty}{Z}^{(i)}(s_i)\,.$$

- Fix $M \in \mathbb{N}$ and consider the partition

$$\mathcal{T}_M := \left\{ \frac{k}{2^M}, \quad k \geq 0 \right\}\,.$$

- Define the following approximations on the partition $\mathcal{T}_M$:

$$\uparrow \overset{\infty}{Z}{}^{(i,M)}_t := \sum_{k=0}^{\infty} \overset{\infty}{Z}{}^{(i)}_{k/2^M} 1_{[\frac{k}{2^M}, \frac{k+1}{2^M})}(t)\,,$$

$$\uparrow \overset{\infty}{\mathcal{Z}}{}^{(M)}(\underline{s}) := \sum_{i=1}^{m} \uparrow \overset{\infty}{Z}{}^{(i,M)}(s_i),$$

The proof

- Set

$$
\begin{aligned}
(t_0, \tau_0, x) &:= (0, 0, x), \\
\overleftarrow{\sigma} &:= (0, \dots, 0), \\
\overrightarrow{\sigma} &:= (\sigma_1^{(1,M)}, \dots, \sigma_1^{(i,M)}, \dots, \sigma_1^{(m,M)})
\end{aligned}
$$

- Solve

$$
\begin{cases}
\dot{\underline{\tau}}((0,0,x);t) = (x + \tilde{\mathcal{Z}})(\underline{\tau}((0,0,x);t)), \\
\underline{\tau}((0,0,x);t_0) = \tau_0 \,,
\end{cases}
$$

for $t \in [0, t_1]$ where

$$
t_1 := \sup\{t > 0 \mid \underline{\tau}((t_0, \tau_0, x); t) \leq \overrightarrow{\sigma}\} \,.
$$

Remark There might be one or more indices i^*, where equality holds. Collect them in a set $I^* \subseteq \{1, \dots, m\}$.

- Update the values

$$\pi_{I^*}\overleftarrow{\sigma} := \pi_{I^*}\overrightarrow{\sigma},$$
$$\pi_{I^*}\overrightarrow{\sigma} := \pi_{I^*}\overrightarrow{\sigma}_{++},$$

where $\overrightarrow{\sigma}_{++}$ contains the next jumps of ${}^{\uparrow}\overset{\infty}{Z}^{(i,M)}$ for all $i \in I^*$ after $\overrightarrow{\sigma}_i$.

- Define

$$\tau_1 := \underline{\tau}((t_0, \tau_0, x); t_1)$$
$$x_1 := x + \Delta^{\uparrow}\overset{\infty}{\mathcal{Z}}^{(M)}(\overleftarrow{\sigma}).$$

- Solve

$$\begin{cases} \dot{\underline{\tau}}((t_1, \tau_1, x_1); t) = (x_1 + \overset{\sim}{\mathcal{Z}})(\underline{\tau}((t_1, \tau_1, x_1); t)), \\ \underline{\tau}((t_1, \tau_1, x_1); t_1) = \tau_1 , \end{cases}$$

for $t \in [t_1, t_2]$ where

$$t_2 := \sup\{t > t_1 \mid \underline{\tau}((t_1, \tau_1, x_1); t) \leq \overrightarrow{\sigma}\}.$$

- Define iteratively, for all $n \geq 1$

$$\begin{aligned} t_{n+1} &:= \sup\{t > 0 \mid \underline{\tau}((t_n, \tau_n, x_n); t) \leq \overrightarrow{\sigma}\}, \\ \tau_{n+1} &:= \underline{\tau}((t_n, \tau_n, x_n); t_{n+1}), \\ x_{n+1} &:= x_n + \Delta^{\uparrow} \overset{\infty}{\mathcal{Z}}{}^{(M)}(\overleftarrow{\sigma}), \end{aligned}$$

where, at each step $\overleftarrow{\sigma}$ and $\overrightarrow{\sigma}$ are updated.

Solution of the approximated problem

Theorem

There exists a solution of

$$\begin{cases} \dot{\underline{\tau}}^{(M)}((t_0,\tau_0,x);t) = (x + \tilde{\mathcal{Z}} + {}^{\uparrow}\tilde{\mathcal{Z}}^{(M)})(\underline{\tau}^{(M)}((t_0,\tau_0,x);t)), \\ \underline{\tau}^{(M)}((t_0,\tau_0,x);t_0) = 0\,. \end{cases}$$

Moreover it holds

$$\lim_{M\to\infty} \underline{\tau}^{(M)}((t_0,\tau_0,x);t) = \underline{\tau}((t_0,\tau_0,x);t)$$

where $\underline{\tau}$ solves

$$\begin{cases} \dot{\underline{\tau}}((t_0,\tau_0,x);t) = (x + \mathcal{Z})(\underline{\tau}((t_0,\tau_0,x);t)), \\ \underline{\tau}((t_0,\tau_0,x);t_0) = \tau_0\,. \end{cases}$$

Thank you for your attention

Bibliography I

Barczy, M., Li, Z., and Pap, G. (2014).
Stochastic differential equation with jumps for multi-type continuous state and continuous time branching processes with immigration.

Caballero, M. E., Pérez Garmendia, J. L., and Uribe Bravo, G. (2013).
A Lamperti-type representation of continuous-state branching processes with immigration.

Duffie, D., Filipović, D., and Schachermayer, W. (2003).
Affine processes and applications in finance.

G., N. and Teichmann, J. (2014).
Pathwise construction of affine processes.

Kallsen, J. (2006).
A didactic note on affine stochastic volatility models.