

Theorem. Let  $0 < \beta < 1$ ,  $\mu^0 > 0$ ,  $\epsilon > 0$ .

If the primal path following method is used with parameter  $\alpha = 1 - \frac{\sqrt{\beta} - \beta}{\sqrt{\beta} + \sqrt{n}}$  and initial primal/dual feasible solution  $x^0 > 0$ ,  $s^0 > 0$ ,  $p^0$  that satisfy

$$\left\| \frac{1}{\mu^0} X_0 S_0 e - e \right\| \leq \beta, \text{ then a primal/dual}$$

feasible solution  $(x^k, p^k, s^k)$  with duality gap

$(s^k)^T x^k \leq \epsilon$  is found after

$$K := \left\lceil \frac{\sqrt{\beta} + \sqrt{n}}{\sqrt{\beta} - \beta} \log \left( \frac{(s^0)^T x^0}{\epsilon} \cdot \frac{1 + \beta}{1 - \beta} \right) \right\rceil = \left\lceil \frac{1}{1 - \alpha} \log \left( \frac{\epsilon_0}{\epsilon} \cdot \frac{1 + \beta}{1 - \beta} \right) \right\rceil$$

iterations.

→ solve an auxiliary LP with  $\mu^0$ ,  $x^0$ ,  $s^0$  such that

$$\left\| \frac{1}{\mu^0} X_0 S_0 e - e \right\| = \frac{1}{4}$$

⇒ # of iterations to reduce the duality gap from

$\epsilon_0 = (s^0)^T x^0$  to  $\epsilon$  is

$$\left\lceil \left(2 + \frac{1}{4}\sqrt{n}\right) \log \left( \frac{\epsilon_0}{\epsilon} \cdot \frac{5}{3} \right) \right\rceil = O \left( \sqrt{n} \log \frac{\epsilon_0}{\epsilon} \right)$$

→ polynomial in  $n$

with  $O((n+n)^3)$  operations in one iteration

$$\parallel \\ O(n^3)$$

## Primal-dual path following algorithm

Idea: Directly try to solve the KKT conditions:

$$Ax(\mu) = b$$

$$x(\mu) \geq 0$$

$$A^T p(\mu) + s(\mu) = c$$

$$s(\mu) \geq 0$$

$$X(\mu)S(\mu)e = e\mu$$

► Newton's method:

Given  $F: \mathbb{R}^r \rightarrow \mathbb{R}^r$ , problem: find a  $z^*$  such that

$$F(z^*) = 0$$

Assume we have an approximation  $z$  for  $z^*$

→ find a direction  $d$  such that  $z+d$  is a better approx.

Taylor series:  $F(z+d) \approx F(z) + J_F(z) \cdot d$

with the Jacobian  $J_F(z) = \left( \frac{\partial}{\partial z_i} F_i(z) \right)_{1 \leq i, j \leq n}$

$$F(z+d) = 0 \Leftrightarrow F(z) + J_F(z) \cdot d = 0$$

► Here:

$$F(x, p, s) = \begin{pmatrix} Ax - b \\ A^T p + s - c \\ XSe - \mu e \end{pmatrix}$$

→ solve the linear system

$$\begin{pmatrix} A & 0 & 0 \\ 0 & A^T & I \\ S & 0 & X \end{pmatrix} \cdot \begin{pmatrix} d_x \\ d_p \\ d_s \end{pmatrix} = -F(x, p, s)$$
  
$$(*) = \begin{pmatrix} 0 \\ 0 \\ \mu e - XSe \end{pmatrix}$$

► Updating the solution and the barrier parameter:

new solution:  $x + \beta_{\text{primal}} d_x$ ,  $p + \beta_{\text{dual}} d_p$ ,  $s + \beta_{\text{dual}} d_s$

$$\text{with } \beta_{\text{primal}} = \min \left\{ 1, \alpha \cdot \min_{\substack{i \\ \text{such that} \\ (d_x)_i < 0}} \left( -\frac{x_i}{(d_x)_i} \right) \right\} \quad (**)$$

$$\beta_{\text{dual}} = \min \left\{ 1, \alpha \cdot \min_{\substack{i \\ \text{such that} \\ (d_s)_i < 0}} \left( -\frac{s_i}{(d_s)_i} \right) \right\}$$

where  $0 < \alpha < 1$ .

$$\bar{\mu} := \rho \cdot \frac{s^T x}{n} \quad \text{with} \quad 0 < \rho \leq 1 \quad (***)$$

► Algorithm:

Input:  $A, b, c, \varepsilon > 0, 0 < \alpha < 1, \mu^0$   
 initial primal & dual feasible solutions  $x^0 > 0, s^0 > 0, p^0$

1.  $k := 0$
2. If  $(s^k)^T x^k < \varepsilon$  then STOP: solution is  $\varepsilon$ -optimal.
3. Let  $\rho^k \in (0, 1]$ , update  $\mu^k$  according to (\*\*\*)  
 Solve the linear system (\*) for  $d_x^k, d_p^k, d_s^k$
4. Find the step lengths  $\beta_{\text{primal}}$  and  $\beta_{\text{dual}}$  according to (\*\*)
5.  $x^{k+1} := x^k + \beta_{\text{primal}} \cdot d_x^k$ ,  
 $p^{k+1} := p^k + \beta_{\text{dual}} \cdot d_p^k, s^{k+1} := s^k + \beta_{\text{dual}} \cdot d_s^k$
6.  $k := k+1$ , go to 2.

Complexity:  $O(\sqrt{n} \log \frac{\epsilon_0}{\epsilon})$  iterations,  
 (even  $O(\log n \log \frac{\epsilon_0}{\epsilon})$  on average?)

Infeasible primal-dual path following method:

Start with  $x^0 > 0$ ,  $s^0 > 0$ ,  $p^0$  that do not necessarily have to be feasible.

Only change: instead of the system (\*) solve:

$$\begin{pmatrix} A & 0 & 0 \\ 0 & A^T & I \\ S_k & 0 & X_k \end{pmatrix} \cdot \begin{pmatrix} d_x^k \\ d_p^k \\ d_s^k \end{pmatrix} = - \begin{pmatrix} Ax^k - b \\ A^T p^k + s^k - c \\ X_k S_k e - \mu^k e \end{pmatrix}.$$


---

Summary:

Interior point methods start with an interior point of the feasible set and try to reach a primal/dual feasible solution pair with duality gap  $< \epsilon$  by proceeding in a certain direction  $d_{\text{method}}$  in every step.

$$d_{\text{affine scaling}} = -X^2 (I - A^T (AX^2 A^T)^{-1} AX^2) c$$

$$d_{\text{path following}} = (I - A^T (AX^2 A^T)^{-1} AX^2) (Xe - \frac{1}{\mu} X^2 c)$$

$$\text{Let } d_{\text{center}} := (I - A^T (AX^2 A^T)^{-1} AX^2) Xe, \text{ then}$$

$$d_{\text{path following}} = d_{\text{center}} + \frac{1}{\mu} d_{\text{affine scaling}}$$

$$d_{\text{potential reduction}} = d_{\text{center}} + \frac{q}{s^T x} d_{\text{affine scaling}}$$

► Comparison with the simplex algorithm:

- in practice interior point methods tend to have a better performance for:

- very large problems

- massively degenerate problems

- interior point methods always find optimal solutions in the interior of the set of all optimal solutions (not a basic optimal solution)

- IP

- sensitive for unbounded sets of optimal solutions

- problems with solving many similar problems