



COMPLEX ANALYSIS I

<http://www3.math.tu-berlin.de/geometrie/Lehre/SS17/ComplexAnalysis/>

EXERCISE SHEET 10

Due before the lecture on Thursday, June 29, 2017.

Exercise 35: Laurent series.

(6 pts)

Let $f : \mathbb{C} \setminus \{1, -2i\} \rightarrow \mathbb{C}$,

$$f(z) = \frac{5}{z^2 + (2i - 1)z - 2i}.$$

1. Determine the Laurent series for f in the annuli

$$\begin{aligned} A_1 &= \{z \in \mathbb{C} \mid 1 < |z| < 2\}, \\ A_2 &= \{z \in \mathbb{C} \mid 0 < |z + 2i| < \sqrt{5}\}, \\ A_3 &= \{z \in \mathbb{C} \mid \sqrt{5} < |z - 1|\}. \end{aligned}$$

2. Compute the following integrals using Part 1:

$$\begin{aligned} &\int_{|z|=\frac{3}{2}} \frac{1}{z^2 + (2i - 1)z - 2i} dz, \\ &\int_{|z+2i|=1} \frac{1}{(z + 2i)^2(z^2 + (2i - 1)z - 2i)} dz. \end{aligned}$$

Exercise 36: Zeros and poles.

(4 pts)

Let f , g and h be holomorphic and let z_0 be a pole of order ℓ of f , a pole of order m of g , and a zero of order n of h . Determine the type of the singularity in z_0 of the maps

$$f + g, \quad f + h, \quad fg, \quad fh, \quad \frac{f}{g}, \quad \frac{f}{h}, \quad \text{and} \quad \frac{h}{f}.$$

Exercise 37: Holomorphic automorphisms of \mathbb{C} .

(6 pts)

- Let $U \subset \mathbb{C}$ be open and let $f : U \rightarrow \mathbb{C}$ be holomorphic and injective. Let z_0 be an isolated singularity of f . Show that z_0 cannot be essential.
- Let $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ be holomorphic and injective. Show that $z_0 = 0$ is either a removable singularity or a pole of first order.
- Let $g : \mathbb{C} \rightarrow \mathbb{C}$ be holomorphic and bijective. Show that $f(z) = g(\frac{1}{z})$ cannot have a removable singularity in $z_0 = 0$. (Thus, it has a pole of first order in z_0 .)
- Determine all biholomorphic functions $g : \mathbb{C} \rightarrow \mathbb{C}$. (These are called holomorphic automorphisms of \mathbb{C} .)¹

¹Hint: Consider the Laurent series of $f(z) = g(\frac{1}{z})$ and use Part 3.