

Differential Geometry II: Analysis and Geometry on Manifolds

Exercise Sheet 2

(Manifolds, diffeomorphisms)

due 06.11.2018

Exercise 1

7 points

Let $\mathbb{R}^n/\mathbb{Z}^n$ denote the quotient space \mathbb{R}^n/\sim with equivalence relation given by

$$x \sim y :\Leftrightarrow x - y \in \mathbb{Z}^n.$$

Let $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^n/\mathbb{Z}^n$, $x \mapsto [x]$ denote the canonical projection. Show:

- π is a *covering map*, i.e. a continuous surjective map such that each point $p \in \mathbb{R}^n/\mathbb{Z}^n$ has a open neighborhood V such that $\pi^{-1}(V)$ is a disjoint union of open sets each of which is mapped by π homeomorphically to V .
- π is an open map.
- $\mathbb{R}^n/\mathbb{Z}^n$ is a manifold of dimension n .
- $\{(\pi|_U)^{-1} \mid U \subset \mathbb{R}^n \text{ open, } \pi|_U: U \rightarrow \pi(U) \text{ bijective}\}$ is a smooth atlas of $\mathbb{R}^n/\mathbb{Z}^n$.

Exercise 2

4 points

Show that the following manifolds are diffeomorphic.

- $\mathbb{R}^2/\mathbb{Z}^2$.
- the product manifold $\mathbb{S}^1 \times \mathbb{S}^1$.
- the torus of revolution as a submanifold of \mathbb{R}^3 :

$$T = \{((R + r \cos \varphi) \cos \theta, (R + r \cos \varphi) \sin \theta, r \sin \varphi) \mid \varphi, \theta \in \mathbb{R}\}.$$

Exercise 3

4 points

Show that the Möbius band (without boundary)

$$M = \{((2 + r \cos \frac{\varphi}{2}) \cos \varphi, (2 + r \cos \frac{\varphi}{2}) \sin \varphi, r \sin \frac{\varphi}{2}) \mid r \in (-\frac{1}{2}, \frac{1}{2}), \varphi \in \mathbb{R}\}$$

is a submanifold of \mathbb{R}^3 . Show further that for each point $p \in \mathbb{R}P^2$ the open set $\mathbb{R}P^2 \setminus \{p\} \subset \mathbb{R}P^2$ is diffeomorphic to M .